

Green approach: A simple one-pot synthesis of pyranopyrazoles scaffold

Ch. Vijaya Saradhi^a, G. Usha Rani^b and P. Rama Krishna Veni^{c*}

^aDepartment of Science and Humanities, KITS Akshar Institute of Technology, NH-16, Yanamadala, Prathipadu Mandalam, Guntur-522019, Andhra Pradesh, India

^bDepartment of Basic Science & Humanities, Vignan's Lara Institute of Technology and Science, Vadlamudi, Guntur-522213, Andhra Pradesh, India

^cDepartment of Applied Sciences & Humanities, Sasi Institute of Technology & Engineering, Tadepalligudem, West Godavari-534101, Andhra Pradesh, India

CHRONICLE

Article history:

Received July 16, 2024

Received in revised form

August 18, 2024

Accepted October 16, 2024

Available online

October 16, 2024

Keywords:

Pyranopyrazole

Zinc acetate. Lewis acid

Four component reaction


Green chemistry and

Catalysis

ABSTRACT

The past decade has witnessed significant progress in synthesizing structurally diverse and biologically relevant pyrano[2,3-c]pyrazole derivatives through the integration of green methodologies. A straightforward and environmentally friendly one-step method has been developed to synthesize divergent pyranopyrazoles in good yields with the aid of zinc acetate as a Lewis acid catalyst in toluene as solvent under reflux conditions.

© 2025 by the authors; licensee Growing Science, Canada.



Graphical Abstract

1. Introduction

The first multicomponent reaction, credited to Strecker and dating to 1850, served as the catalyst for a string of reactions that have been documented in literature.¹ Multicomponent reactions (MCRs) are a unique class of synthetically useful organic reactions in which one or more different starting ingredients are combined in one pot to form a final product.²⁻⁴ These reactions have been carefully used in a number of synthetic transformations, where traditional techniques typically demand for a lengthy process with numerous phases. High yields, atom-/step economy, shortened reaction times, environmental friendliness, and a useful tool for building a library of novel chemical entities (NCEs) are all benefits of the MCR technique, which is particularly useful in the drug development process.⁵⁻⁹ In heterocyclic chemistry, the creation of highly convergent syntheses has been a constant since heterocyclic scaffolds frequently consist of more than two building units.¹⁰ Heterocycles' triumphant journey toward functional molecules has been closely linked to their synthesis by multicomponent reactions (MCRs) from the early days of organic chemistry.


* Corresponding author

E-mail address pramaky2023@gmail.com (P. R. K. Veni)

© 2025 by the authors; licensee Growing Science, Canada

doi: 10.5267/j.ccl.2024.10.008

As a subgroup of heterocycles, pyranopyrazoles have attracted a lot of interest because of their various structural roles and biological activity.¹¹⁻¹⁴ The four different isomer configurations of these compounds—pyrano[2,3-*c*]pyrazole, pyrano[3,2-*c*]pyrazole, pyrano[3,4-*c*]pyrazole, and pyrano[4,3-*c*]pyrazole—are made up of fused pyran and pyrazole rings (Fig. 1).



But because this isomer has biological importance, pyrano[2,3-*c*]pyrazoles are the ones that have been studied the most. These substances have demonstrated encouraging antiviral, antibacterial, anticancer, and anti-inflammatory qualities. Furthermore, they demonstrate the capacity to conceivably suppress the function of the human Chk1 kinase enzyme (Figure 2).¹⁵⁻¹⁶ Because of their structural variety, it is possible to alter different parts of the molecule to affect their activity, which creates opportunities for research on the link between structure and activity. In the one-pot multicomponent reaction of pyranopyrazoles, the overall reaction scheme usually consists of an aldehyde, malononitrile, hydrazine hydrate, β -ketoester/ethyl acetoacetate, and a suitable catalyst or promoter. Condensation, cyclization, and subsequent rearrangement are some of the sequential transformations that the reaction goes through to produce the pyranopyrazole product. Undoubtedly, a great deal of study has gone into the synthesis of pyrano[2,3-*c*]pyrazole, leading to the discovery of a variety of techniques and synthetic pathways. Pyranopyrazole synthesis has been approached from different angles using catalysts like PhCO_2Na , thiamine hydrochloride, taurine, Fe/Cu nanocomposites, ethylene glycol (E-G), Fe_3O_4 @chitosan-tannic acid bionanocomposite, piperidine, SnCl_2 , dodecylbenzene sulfonic acid (DBSA), PTSA, ionic liquids, citric acid, β -cyclodextrin, NH_4Cl , ZrO_2 -NPs, PS-PTSA, nano-eggshell/Ti(IV), thiamine hydrochloride, heteropolyacid, PhCO_2Na , chitosan hydrogel, taurine, Fe/Cu nanocomposites, ethylene glycol (E-G), Fe_3O_4 @chitosan-tannic acid bionanocomposite, etc.¹⁷⁻⁴⁶ However, the majority of the material now in publication concentrates on attaining high yields and product diversity, frequently overshadowing the crucial significance of sustainability.




Fig. 2. Some biologically active pyrano[2,3-*c*]pyrazoles

Of the several zinc salts that are available, zinc acetate is one of the inexpensive, more easily obtained, and less toxic Lewis acids. Because of its peculiar physical and chemical characteristics, which show that it can be helpful in enabling a

broad range of synthetic transformations in both organic synthesis and catalysis, it is known as a multifunctional catalyst.⁴⁷⁻⁵⁴ Zinc acetate is affordable, easy to produce, and stable in the presence of oxygen and moisture when used in laboratory conditions. As a follow-up to earlier studies and a current addition to green synthetic methods, we report a one-pot multicomponent synthesis of pyranopyrazole. Following a comprehensive analysis of all pertinent published data, we came to the conclusion that $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$ was not involved in the relevant reaction.

2. Results and Discussion

2.1 Effect of zinc salts, solvents and temperature


Later, various zinc catalysts were examined under controlled circumstances to determine how well they supported the control reaction on benzaldehyde, hydrazine hydrate, malanonitrile and ethyl acetoacetate (**Table 1**). $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$ (10 mol%) was found to be the most effective catalyst when compared to the other catalysts tested (ZnCl_2 , ZnO , $\text{Zn}(\text{NO}_3)_2$, ZnCO_3 and ZnSO_4) in terms of reaction time and yield at reflux temperature in toluene as solvent. The control reaction was determined to be viable at 110 °C. The related product was produced in minimal or small amounts when the temperature was gradually lowered till ambient temperature. The reaction can produce the best results at 5 mol%, according to studies of catalyst loading (5, 7.5, and 10 mol%). Higher catalyst concentrations (12.5 and 15 mol%) had no effect on the yield either up or down.

Table 1. Effect of available zinc salts tested

Entry	Zinc salt	Reaction time (h)	Isolated yield (%) ^a
1	ZnCl_2	4.5	64
2	ZnO	24	7
3	$\text{Zn}(\text{NO}_3)_2$	48	11
4	ZnCO_3	48	NR
5	ZnSO_4	24	NR
5	$\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$	2	89

Toluene emerged as the superior solvent under reflux conditions proved to be the most effective reaction condition (**Table 2**, Entry 1), surpassing other less efficient solvents (Table 2, entries 2-6). Non polar solvents (entries 1 and 6) proved to be better compared to that of polar solvents (entries 2-5) as evident from Table 2. For this reason, we further prepared derivatives from the above standard reaction condition (**Figure 3**). The results revealed that aldehydes with electron-withdrawing groups or electron-donating groups have similar effect. Important to note is functional group ($-\text{NO}_2$) be at *ortho*-, *meta*-, or *para*- position has no steric effect to form the corresponding pyranopyrazoles (**5f**, **5g** and **5h**, Figure 3). Heteroaromatic aldehyde such as furfural gave the product **5i** in good yield. However, aliphatic aldehyde (*n*-butanal) reacted sluggishly to give the corresponding product **5j**.

Table 2. Optimization of the solvent and catalyst loading



Entry	Solvent	Temperature (°C)	Time (h)	Yield (%) ^{a,b}
1	PhCH_3	Room temperature 110 °C	8 2	23 89
2	CH_3CN	Reflux	6	18
3	EtOAc	Reflux	4	43
4	EtOH	Reflux	8	28
5	CHCl_3	Reflux	6	30
6	CCl_4	Reflux	6	78
7	Without Solvent	Room temperature	12	NR

^aYields refer to pure isolated product

^b $\text{Zn}(\text{OAc})_2 \cdot 2\text{H}_2\text{O}$ as Lewis catalyst loading: (89%, 10 mol%); (82%, 7.5 mol%) and (80%, 5 mol%)




Fig. 3. Synthesized dihydropyranopyrazole[2,3-*c*]pyrazoles

3. Experimental

3.1 Material and Methods

All reagents and chemicals were of analytical grade and used without further purification. The progress of the reaction was monitored on TLC. All the products were confirmed by comparing their melting points, IR, and ^1H NMR data with literature data.

General procedure for the synthesis of substituted pyranopyrazole (5a-j)

To a round bottom flask substituted aromatic aldehyde (1mmol), malononitrile (1mmol), ethyl acetoacetate (1mmol), and hydrazine hydrate (1 mmol) in toluene under reflux conditions for the time as specified in **Figure 3**. The advancement of the reaction was tracked through TLC (10% ethyl acetate: *n*-hexane). The resulting product was combined with water and ethyl acetate (1:4). The combined solvent extracts underwent vacuum concentration. Subsequently, the compounds underwent recrystallization in ethanol to yield the purified product (5a-j).

Spectral data of synthesized compounds

6-Amino-3-methyl-4-phenyl-1,4-dihydropyranopyrazole-5-carbonitrile (5a):

Yellow crystalline solid. **m.p.** 210-212 $^{\circ}\text{C}$. **IR (KBr) cm^{-1}** - 3477 and 3231 (NH_2), 3121 (NH), 2191 (CN), 1639, 1559. **^1H NMR-(400 MHz, DMSO *d*₆)** - δ ppm 12.00 (1H, s, NH), 7.30 (2H, t, $J = 7.56$), 7.22 (3H, m, $J = 7.28$ and 11.24), 6.71 (2H, s, $-\text{NH}_2$), 4.50 (1H, s, C-4), 1.79 (3H, s, CH_3). **^{13}C NMR- (100 MHz, DMSO *d*₆)** - 161.35, 155.26, 144.93, 136.04, 128.91, 127.94, 127.20, 121.24, 98.13, 57.71, 36.73, 10.20. **MS (EI): m/z Calcd.** for $\text{C}_{14}\text{H}_{12}\text{N}_4\text{O} = 252.10$, Found = 253.15 [$\text{M}^+ + 1$].

6-Amino-1,4-dihydro-3-methyl-4-*p*-tolylpyranopyrazole-5-carbonitrile (5b):

White crystalline solid. **m.p.** 207-208 $^{\circ}\text{C}$. **IR (KBr) cm^{-1}** - 3406 and 3315 (NH_2), 3188 (NH), 2191 (CN), 1646 (C=N), 1600 (C=C). **^1H NMR-(400 MHz, DMSO *d*₆)** - δ ppm 12.19 (1H, s, NH), 8.21 (2H, m, $J = 8.8$ Hz, Ar), 7.47 (2H, m, $J = 8.8$ Hz, Ar), 7.03 (2H, s, $-\text{NH}_2$), 4.82 (1H, s, C-4), 3.73 (s, 3H, OCH_3), 1.78 (3H, s, CH_3). **^{13}C NMR- (100 MHz, DMSO *d*₆)** - δ ppm 161.06, 154.58, 152.02, 146.30, 135.84, 128.51, 128.77, 123.83, 120.45, 96.48, 55.80, 35.80, 9.66. **MS: m/z Calcd.** for $\text{C}_{15}\text{H}_{14}\text{N}_4\text{O} = 266.12$, Found = 267.14 [$\text{M}^+ + 1$].

6-Amino-1,4-dihydro-4-(4-methoxyphenyl)-3-methylpyranopyrazole-5-carbonitrile (5c):

Light yellow crystalline solid. **m.p.** 209-210 $^{\circ}\text{C}$. **IR (KBr) cm^{-1}** - 3483 and 3249 (NH_2), 3122 (NH), 2190 (CN), 1643 (C=N), 1600 (C=C). **^1H NMR-(400 MHz, DMSO *d*₆)** - δ ppm 12.08 (1H, s, NH), 7.10 (2H, m, $J = 8$ Hz, Ar), 6.89 (2H, m, $J = 4$ Hz, Ar), 6.82 (2H, s, $-\text{NH}_2$), 4.55 (1H, s, C-4), 3.74 (3H, s, OCH_3), 1.80 (3H, s, CH_3). **^{13}C NMR- (100 MHz, DMSO *d*₆)** - δ ppm 161.16, 158.45, 155.24, 136.96, 136.02, 128.96, 121.28, 114.25, 98.37, 58.15, 55.48, 35.93, 10.21. **MS: m/z Calcd.** for $\text{C}_{15}\text{H}_{14}\text{N}_4\text{O}_2 = 282.12$, Found = 283.16 [$\text{M}^+ + 1$].

6-Amino-4-(4-chlorophenyl)-1,4-dihydro-3-methylpyrano[2,3-c]pyrazole-5-carbonitrile (5d):

Yellow crystalline solid. **m.p.** 228-230 °C. **IR (KBr) cm^{-1}** - 3373 and 3311 (NH_2), 3171 (NH), 2193 (CN), 1627, 1581, 1372, 1255, 879. **^1H NMR-(400 MHz, DMSO d_6)** - δ ppm 12.12 (1H, s, NH), 7.38 (2H, m, $J = 11.2$ Hz, Ar), 7.21 (2H, m, $J = 11.2$ Hz, Ar), 6.90 (2H, s, NH_2), 4.63 (1H, s, C-4), 1.79 (3H, s, CH_3). **^{13}C NMR-(100 MHz, DMSO d_6)** - δ ppm 160.84, 154.61, 143.38, 135.68, 132.49, 131.19, 129.96, 128.39, 120.62, 97.12, 56.68, 18.45, 9.66. **MS:** m/z Calcd. for $\text{C}_{14}\text{H}_{11}\text{ClN}_4\text{O}$ = 286.06, Found = 287.10 [$\text{M}^+ + 1$].

6-Amino-4-(2,4-dichlorophenyl)-3-methyl-1,4-dihydropyrano[2,3-c]pyrazole-5-carbonitrile (5e):

White crystalline solid. **m.p.** 223-225 °C; **IR (KBr) cm^{-1}** - 3478 and 3246 (NH_2), 3117 (NH), 2190 (CN), 1640, 1504, 1408, 1100, 1052, 866, 741. **^1H NMR-(400 MHz, DMSO d_6)** - δ ppm 11.87 (s, 1H, NH), 7.70 (d, 2H, $J = 8.4$ Hz, ArH), 7.62 (d, 2H, $J = 8.4$ Hz, ArH), 7.13 (s, 2H, NH_2), 7.02 (s, 1H, ArH), 4.58 (s, 1H, C-4), 1.74 (s, 3H, CH_3). **^{13}C NMR- (100 MHz, DMSO d_6)** - δ ppm 161.8, 155.4, 140.6, 135.9, 133.3, 132.6, 129.3, 128.5, 120.7, 96.8, 55.7, 33.6, 10.0. **MS:** Exact mass: (M^+): calcd. 320.0232; found 320.0236.

6-Amino-1,4-dihydro-3-methyl-4-(3-nitrophenyl)pyrano[2,3-c]pyrazole-5-carbonitrile (5f):

Brown crystalline solid. **m.p.** 214-215 °C. **IR (KBr) cm^{-1}** - 3433 and 3334 (NH_2), 3202 (NH), 2189 (CN), 1668, 1529, 1419, 1358. **^1H NMR-(400 MHz, DMSO d_6)** - δ ppm 12.19 (1H, s, NH), 8.13 (1H, m, $J = 10.4$ Hz, Ar), 8.02 (1H, m, $J = 1.2$ Hz, Ar), 7.63 (2H, s, Ar), 7.03 (1H, s, NH_2), 4.87 (1H, s, C-4), 1.80 (3H, s, CH_3). **^{13}C NMR- (100 MHz, DMSO d_6)** - δ ppm 161.07, 154.61, 147.80, 146.74, 135.88, 134.34, 130.19, 121.94, 121.77, 120.49, 96.59, 56.04, 35.55, 9.68. **MS:** m/z Calcd. for $\text{C}_{14}\text{H}_{11}\text{N}_5\text{O}_3$ = 297.09, Found = 298.13 [$\text{M}^+ + 1$].

6-Amino-3-methyl-4-(2-nitrophenyl)-1,4-dihydropyrano[2,3-c]pyrazole-5-carbonitrile (5g): Light yellow crystalline solid. **m.p.** 214-215 °C. **IR (KBr) cm^{-1}** - 3420 and 3351 (NH_2), 3167 (NH), 2206 (CN). **^1H NMR-(400 MHz, DMSO d_6)** - δ ppm 12.2 (s, 1H, NH), 7.88 (d, $J = 8.0$ Hz, 1H, HAr), 7.70 (t, $J = 7.5$ Hz, 1H, HAr), 7.53 (t, $J = 7.0$ Hz, 1H, HAr), 7.35 (d, $J = 8.0$ Hz, 1H, HAr), 7.08 (s, 2H, NH_2), 5.12 (s, 1H, C-4), 1.80 (s, 3H, CH_3).

6-Amino-1,4-dihydro-3-methyl-4-(4-nitrophenyl)pyrano[2,3-c]pyrazole-5-carbonitrile (5h)

Yellow crystalline solid. **m.p.** 249-250 °C. **IR (KBr) cm^{-1}** - 3310 and 3165 (NH_2), 2900 (NH), 2185 (CN), 1642, 1596. **^1H NMR-(400 MHz, DMSO d_6)** - δ ppm 12.19 (1H, s, NH), 8.21 (2H, m, $J = 8.8$ Hz, Ar), 7.47 (2H, m, $J = 8$ Hz, Ar), 7.10 (2H, s, NH_2), 4.82 (1H, s, C-4), 1.79 (3H, s, CH_3). **^{13}C NMR- (100 MHz, DMSO d_6)** - δ ppm 161.06, 154.58, 152.01, 146.30, 135.86, 129.52, 128.77, 124.07, 120.45, 96.48, 55.80, 35.79, 9.66. **MS:** m/z Calcd. for $\text{C}_{14}\text{H}_{11}\text{N}_5\text{O}_3$ = 297.09, Found = 298.14 [$\text{M}^+ + 1$].

6-Amino-4-(furan-2-yl)-3-methyl-2,4-dihydropyrano[2,3-c]pyrazole-5-carbonitrile (5i)

Grey solid. **m.p.** 215-216 °C. **IR (KBr) cm^{-1}** - 3470 and 3401 (NH_2), 3120 (NH), 2189 (CN). **^1H NMR-(400 MHz, DMSO d_6)** - δ ppm 12.25 (s, 1H, NH), 7.49 (s, 1H, CH, Ar), 6.33 (s, 1H, CH, Ar), 6.17 (s, 1H, CH), 4.75 (s, 1H, C-4), 1.96 (s, 3H, CH_3) ppm; **^{13}C NMR- (100 MHz, DMSO d_6)** - δ ppm 162.0, 156.22, 155.32, 142.81, 136.35, 121.35, 110.77, 106.17, 95.62, 54.43, 25.50, 10.11.

6-Amino-4-(4-isopropylphenyl)-3-methyl-1,4-dihydropyrano[2,3-c]pyrazole-5-carbonitrile (5j): White solid, **m.p.** 239-241 °C, **IR (KBr, cm^{-1})**- 3494 and 3233 (NH_2), 2961 (NH), 2196 (CN), 1613, 1597, 1490, 1398, 1053. **^1H NMR-(400 MHz, DMSO d_6)** - δ 12.06 (s, 1H, NH), 7.16 (d, $J = 7.9$ Hz, 2H, Ar), 7.04 (d, $J = 7.8$ Hz, 2H, Ar), 6.83 (s, 2H), 4.52 (s, 1H, C-4), 2.81 (q, $J = 6.9$ Hz, 1H), 1.77 (s, 3H), 1.16 (d, $J = 6.9$ Hz, 6H).

Supporting Information file is available

4. Conclusion

In summary, we successfully implemented a one-pot multicomponent synthesis of pyranopyrazole using an environmentally friendly synthetic approach with inexpensive and less toxic zinc acetate using as a Lewis acid catalyst. Aromatic compounds both with electron donating and withdrawing functionalities reacted in a similar fashion. Whereas, aliphatic aldehydes proved to be less effective compared to that of reactivity of aromatic compounds. This method utilizes toluene as a solvent and offers benefits such as a shorter reaction duration, straightforward workup, and atom economy. The established protocol is robust and may be used in a variety of organic transformations to reach the desired architecture's complexity in a single pot.

Conflict of Interests

The authors declare that there is no conflict of interest.

References

1. Strecker A. (1850) Ueber die künstliche Bildung der Milchsäure und einen neuen, dem Glycocoll homologen Körper. *Liebigs Ann. Chem.*, 75(1) 27-45. <https://doi.org/10.1002/jlac.18500750103>
2. Tryfon Z-T., Ajay L. C., and Alexander D. (2015) Multicomponent reactions, union of MCRs and beyond. *The Chemical Record* 15(5) 981-996. <https://doi.org/10.1002/tcr.201500201>
3. Zarganes-Tzitzikas T., and Dömling A. (2014) Modern multicomponent reactions for better drug syntheses. *Org. Chem. Front.*, 1 834-837. <https://doi.org/10.1039/C4QO00088A>
4. Cores Á., Clerigué J., Orocio-Rodríguez E., and Menéndez J. C. (2022) Multicomponent Reactions for the synthesis of active pharmaceutical ingredients. *Pharmaceuticals* 15 1009. <https://doi.org/10.3390/ph15081009>
5. Younus H. A., Al-Rashida M., Hameed A., Uroos M., Salar U., Rana S., and Khan K. M. (2020) Multicomponent reactions (MCR) in medicinal chemistry: a patent review (2010-2020). *Expert Opinion on Therapeutic Patents* 31(3) 267-289. <https://doi.org/10.1080/13543776.2021.1858797>
6. Graziano G., Stefanachi A., Contino M., Prieto-Díaz R., Ligresti A., Kumar P., Scilimati A., Sotelo E., and Leonetti F. (2023) Multicomponent reaction-assisted drug discovery: A time- and cost-effective green approach speeding up identification and optimization of anticancer drugs. *Int. J. Mol. Sci.*, 24 6581. <https://doi.org/10.3390/ijms24076581>
7. Graebin C. S., Ribeiro F. V., Rogério K. R., and Kümmerle A. E. (2019) Multicomponent reactions for the synthesis of bioactive compounds: A review. *Curr Org. Synth.*, 16(6) 855-899. <https://doi.org/10.2174/1570179416666190718153703>
8. John S. E., Shivani G., and Nagula S. (2021) Recent advances in multi-component reactions and their mechanistic insights: a triennium review. *Org. Chem. Front.*, 8 4237-4287. <https://doi.org/10.1039/D0QO01480J>
9. Godsi M. Z., Moradi R., and Mahammadkhani L. (2019) Application of multicomponent reactions in the total synthesis of natural peptides. *Arkivoc* part i 18-40. <https://doi.org/10.24820/ark.5550190.p010.779>
10. Multicomponent reactions in the synthesis of heterocycles. (2017) *Chem. Heterocycl. Comp.*, 53 381. <https://doi.org/10.1007/s10593-017-2064-2>
11. Yadav A. R., Katariya A. P., Kanagare A. B., Patil P. D. J., Tagad C. K., Dake S. A., and Nagwade P. A. (2024) Review on advancements of pyranopyrazole: synthetic routes and their medicinal applications. *Mol. Divers.*, <https://doi.org/10.1007/s11030-023-10757-w>
12. Mamaghani M., and Nia, R. H. (2019) A review on the recent multicomponent synthesis of pyranopyrazoles. *Polycyclic Aromatic Compounds* 41(2) 223-291. <https://doi.org/10.1080/10406638.2019.1584576>
13. Ganta R. K., Kerru N., Maddila S., and Jonnalagadda S. B. (2021) Advances in pyranopyrazole scaffolds' syntheses using sustainable catalysts-A review. *Molecules* 26(11) 3270. <https://doi.org/10.3390/molecules26113270>
14. Nagarajan A. S., and Reddy B. S. R. (2009) Synthesis of substituted pyranopyrazoles under neat conditions via a multicomponent reaction. *Synlett* 12 2002-2004. <https://doi.org/10.1055/s-0029-1217526>
15. Biswas S. K., and Das D. (2022) One-pot synthesis of pyrano[2,3-*c*]pyrazole derivatives via multicomponent reactions (MCRs) and their applications in medicinal chemistry. *Mini Rev. Org. Chem.*, 19(5) 552-568. <https://doi.org/10.2174/1570193X19666211220141622>
16. Ahmad A., Rao S., and Shetty N. S. (2023) Green multicomponent synthesis of pyrano[2,3-*c*]pyrazole derivatives: current insights and future directions. *RSC Adv.*, 13(41) 28798-28833. <https://doi.org/10.1039/d3ra05570a>
17. Sikandar S., and Zahoor A. F. (2021) Synthesis of pyrano[2,3-*c*]pyrazoles: A review. *J. Heterocycl. Chem.*, 58(3) 685-705. <https://doi.org/10.1002/jhet.4191>
18. Letcy V. T., Savitha D. P., Deepa S., and Sreekumar K. (2021) Sustainable multicomponent one pot synthesis of pyranopyrazole derivatives in the presence of Lactic acid: Urea: NH₄Cl. *Current Research in Green and Sustainable Chemistry* 4 100194. <https://doi.org/10.1016/j.crgsc.2021.100194>
19. Farooq S., and Ngaini Z. (2024) Recent synthesis of mono- & bis-pyranopyrazole derivatives. *ChemistrySelect* 9(8) e202400028. <https://doi.org/10.1002/slct.202400028>
20. Zolfigol M. A., Tavasoli M., Moosavi-Zare A. R., Moosavi P., Kruger H. G., Shiri, M., and Khakyzadeh V. (2013) Synthesis of pyranopyrazoles using isonicotinic acid as a dual and biological organocatalyst. *RSC Adv.*, 3 25681-25685. <https://doi.org/10.1039/C3RA45289A>
21. Abhinav B., Dinesh C., Alpna A., Pratyush K., and Shrikant M. (2020) Green synthesis of pyranopyrazole using microwave assisted techniques. *GSC Biological and Pharmaceutical Sciences* 10(02) 111-119. <https://doi.org/10.30574/gscbps.2020.10.2.0026>
22. Nguyen H. T., Truong M-N. H., Le T. V., Vo N. T., Nguyen H. D., and Phuong H. T. (2022) A new pathway for the preparation of pyrano[2,3-*c*]pyrazoles and molecular docking as inhibitors of p38 MAP kinase. *ACS Omega* 7(20) 17432-17443. <https://pubs.acs.org/doi/10.1021/acsomega.2c01814>
23. Hanuman D. B., Dattatraya N. P., Ajit K. D., and Rajita D. I. (2024) A facile single-pot synthesis of pyranopyrazoles scaffold as a green approach. *Rasayan J. Chem.*, 17(3) 855-860. <http://doi.org/10.31788/RJC.2024.1738818>

24. Noory Fajer A., Khabt Aboud H., Al-Bahrani H. A., and Kazemi M. (2023) Recent advances on multicomponent synthesis of pyranopyrazoles using magnetically recoverable nanocatalysts. *Polycyclic Aromatic Compounds* 44(7) 4932-4978. <https://doi.org/10.1080/10406638.2023.2255723>
25. Shrivatas P., Pandey R., Zodape S., Wankhade A., and Pratap U. (2020) Green synthesis of pyranopyrazoles via biocatalytic one-pot Knoevenagel condensation–Michael-type addition–heterocyclization cascade in non-aqueous media. *Res. Chem. Intermed.*, 46 2805-2816. <https://doi.org/10.1007/s11164-020-04122-x>
26. Cahyana A. H., and Mufidah A. (2021) Synthesis of pyranopyrazole derivative compounds with nano-Fe₃O₄ catalyst. *AIP Conf. Proc.* 2374 040001. <https://doi.org/10.1063/5.0060453>
27. Khan Md. M., Singh B., Arif A., Saigal., and Sahoo S. C. (2024) Organobase catalyzed synthesis of pyranopyrazoles with X-ray crystallography, docking and ADME studies. *Tetrahedron Green Chem.*, 4 100050. <https://doi.org/10.1016/j.tgchem.2024.100050>
28. Dehghani T. A., Mirjalili B. B. F., Bamoniri A., and Salehi N. (2021) Rapid four-component synthesis of dihydropyrano[2,3-*c*]pyrazoles using nano-eggshell/Ti(IV) as a highly compatible natural based catalyst. *BMC Chemistry*, 15 6. <https://doi.org/10.1186/s13065-021-00734-5>
29. Aliabadi R. S., and Mahmoodi N. O. (2016) Green and efficient synthesis of pyranopyrazoles using [bmim][OH⁻] as an ionic liquid catalyst in water under microwave irradiation and investigation of their antioxidant activity. *RSC Adv.*, 6 85877-85884. <https://doi.org/10.1039/C6RA17594E>
30. Chavan H. V., Babar, S. B., Hoval R. U., and Bandgar, B. P. (2011) Rapid one-pot, four component synthesis of pyranopyrazoles using heteropolyacid under solvent-free condition. *Bull. Korean Chem. Soc.*, 32(11) 3963-3966. <https://doi.org/10.5012/bkcs.2011.32.11.3963>
31. Kiyani H., Samimi H. A., Ghorbani F., and Esmaili S. (2013) One-pot, four-component synthesis of pyrano[2,3-*c*]pyrazoles catalyzed by sodium benzoate in aqueous medium. *Curr. Chem. Lett.*, 2 197-206. <https://doi.org/10.5267/j.ccl.2013.07.002>
32. Patil K., and Helavi V. (2018) Synthesis of pyranopyrazoles by using chitosan hydrogel as a green and recyclable catalyst. *Asian J. Research Chem.*, 11(2) 477-484. <https://doi.org/10.5958/0974-4150.2018.00087.1>
33. Sharma A., Chowdhury R., Dash S., Pallavi B., and Shukla, P. (2016) Fast microwave assisted synthesis of pyranopyrazole derivatives as new anticancer agents. *Current Microwave Chemistry* 3(1) 78-84. <https://doi.org/10.2174/2213335602666150116233238>
34. Neysi M., and Elhamifar D. (2023) H77j8 Yolk-shell structured magnetic mesoporous organosilica supported ionic liquid/Cu complex: an efficient nanocatalyst for the green synthesis of pyranopyrazoles. *Front. Chem.*, 11 1235415. <https://doi.org/10.3389/fchem.2023.1235415>
35. Mohamadpour F. (2020) Catalyst-free green synthesis of dihydropyrano[2,3-*c*]pyrazole scaffolds assisted by ethylene glycol (E-G) as a reusable and biodegradable solvent medium. *J. Chem. Sci.*, 132 72. <https://doi.org/10.1007/s12039-020-01775-4>
36. Kamalzare M., Ahghari M. R., Bayat M., and Maleki A. (2021) Fe₃O₄@chitosan-tannic acid bionanocomposite as a novel nanocatalyst for the synthesis of pyranopyrazoles. *Sci. Rep.*, 11 20021. <https://doi.org/10.1038/s41598-021-99121-2>
37. Al-Amiery A. A., Al-Bayati R. I., Saed F.M., Ali W. B., Kadhum A. A. H., and Mohamad A. B. (2012) Novel pyranopyrazoles: Synthesis and theoretical studies. *Molecules* 17 10377-10389. <https://doi.org/10.3390/molecules170910377>
38. Khazaal S., and Ibraheem S. Y. (2022) One-pot synthesis of new pyranopyrazoles via domino multicomponent reaction. *Egyptian Journal of Chemistry* 65(10) 259-265. <https://doi.org/10.21608/ejchem.2022.115337.5232>
39. Sharma A., Pallavi B., Singh R. P., Jha P. N., and Shukla P. (2015) Novel grinding synthesis of pyranopyrazole analogues and their evaluation as antimicrobial agents. *Heterocycles* 91(8) 1615-1627. <https://doi.org/10.3987/COM-15-13222>
40. Mali G., Shaikh B. A., Garg S., Kumar A., Bhattacharyya S., Erande R. D., and Chate A. V. (2021) Design, synthesis, and biological evaluation of densely substituted dihydropyrano[2,3-*c*]pyrazoles via a taurine-catalyzed green multicomponent approach. *ACS Omega* 6(45) 30734-30742. <https://doi.org/10.1021/acsomega.1c04773>
41. Beiranvand M., and Habibi D. (2022) Design, preparation and application of the semicarbazide-pyridoyl-sulfonic acid-based nanocatalyst for the synthesis of pyranopyrazoles. *Sci. Rep.*, 12 14347. <https://doi.org/10.1038/s41598-022-18651-5>
42. Barhoumi A., Ourhiss N., Belghiti M. E., Chafi M., Syed A., Eswaramoorthy R., Verma M., Zeroual A., Zawadziska K., and Jasiński R. (2023) 3-Difluoromethyl-5-carbomethoxy-2,4-pyrazole: Molecular mechanism of the formation and molecular docking study. *Curr. Chem. Lett.*, 12 472-488. [10.5267/j.ccl.2023.3.008](https://doi.org/10.5267/j.ccl.2023.3.008)
43. Fryźlewicz A., Kačka-Zych A., Demchuk O. M., Mirosław B., Woliński P., and Jasiński R. (2021) Green synthesis of nitrocyclopropane-type precursors of inhibitors for the maturation of fruits and vegetables via domino reactions of diazoalkanes with 2-nitroprop-1-ene. *J. Clean. Prod.*, 292 126079. <https://doi.org/10.1016/j.jclepro.2021.126079>
44. Kula K., Dobosz J., Jasiński R., Kačka-Zych A., Łapczuk-Krygier A., Mirosław B., and Demchuk O. M. (2020) [3+2] Cycloaddition of diaryldiazomethanes with (*E*)-3,3,3-trichloro-1-nitroprop-1-ene: An experimental, theoretical and structural study. *J. Mol. Struct.*, 1203 127473. <https://doi.org/10.1016/j.molstruc.2019.127473>
45. Kula K., Łapczuk A., Sadowski M., Kras J., Zawadziska K., Demchuk O. M., Gaurav G. K., Wróblewska A., and Jasiński R. (2022) On the question of the formation of nitro-functionalized 2,4-pyrazole analogs on the basis of

- nitrylimine molecular systems and 3,3,3-trichloro-1-nitroprop-1-ene. *Molecules* 27 8409. <https://doi.org/10.3390/molecules27238409>
46. Fryźlewicz A., Olszewska A., Zawadzińska K., Woliński P., Kula K., Kačka-Zych A., Łapczuk-Krygier A., and Jasiński R. (2022) On the mechanism of the synthesis of nitrofunctionalised Δ^2 -pyrazolines via [3+2] Cycloaddition reactions between α -EWG-activated nitroethenes and nitrylimine TAC Systems. *Organics* 3 59-76. <https://doi.org/10.3390/org3010004>
 47. Alam M. M., Varala R., and Seema V. (2024) Zinc acetate in organic synthesis and catalysis: A review. *Min. Rev. Org. Chem.*, 21 555-587. <https://doi.org/10.2174/1570193X20666230507213511>
 48. Babu H. B., Varala R., and Alam M. M. (2022) Zn(OAc)₂·2H₂O-Catalyzed Betti base synthesis under solvent free conditions. *Lett. Org. Chem.*, 19, 14-18. <https://doi.org/10.2174/1570178618666210616155257>
 49. Reddy V. V. R., Saritha B., Ramu R., Varala R., and Jayashree, A. (2014) Zn(OAc)₂·2H₂O-catalyzed one-pot efficient synthesis of aminonitriles. *Asian J. Chem.*, 26 7439-7442. <https://doi.org/10.14233/ajchem.2014.17180>
 50. Ramu E., Varala R., Sreelatha N., and Adapa S. R. (2007) Zn(OAc)₂·2H₂O: A versatile catalyst for the one-pot synthesis of propargylamines. *Tetrahedron Lett.*, 48 7184-7190. <https://doi.org/10.1016/j.tetlet.2007.07.196>
 51. Kokane B. D., Varala R., and Patil S. G. (2022) Zn(OAc)₂·2H₂O: An efficient catalyst for the one-pot synthesis of 2-substituted benzothiazoles. *Org. Commun.*, 15 378-385. <https://doi.org/10.25135/acg.oc.140.2210.2618>
 52. Pulle J. S., Totawar P. R., and Varala R. (2023) Zn(OAc)₂·2H₂O-Catalyzed green synthesis of substituted 1-amido/thioamidoalkyl-2-naphthols. *Rev. Roum. Chim.*, 68 75-83. <https://doi.org/10.33224/rch.2023.68.1-2.07>
 53. Chinta B., Satyadev T. N. V. S. S., and Adilakshmi G. V. (2023) Zn(OAc)₂·2H₂O-catalyzed one-pot synthesis of divergently substituted imidazoles. *Curr. Chem. Lett.*, 12 175-184. <https://doi.org/10.5267/j.ccl.2023.3.005>
 54. Veni P. R. K., Saradhi Ch. V., and Rani G. U. (2024) Zn(OAc)₂·2H₂O-catalyzed synthesis of chromeno[2,3-*d*]pyrimidinones under solvent-free conditions. *Curr. Chem. Lett.*, 13 677-682. <https://doi.org/10.5267/j.ccl.2024.5.003>



© 2025 by the authors; licensee Growing Science, Canada. This is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).