

On the question of the correlation between kinetic Dimroth parameters and global electron density transfer in [4+2]- π -electron cycloaddition reactions

Karolina Kula^{a*}, Agnieszka Łapczuk^a, Przemysław Woliński^a and Mikołaj Sadowski^a

^aDepartment of Organic Chemistry and Technology, Cracow University of Technology, Warszawska 24, 31-155 Cracow, Poland

CHRONICLE

Article history:

Received October 12, 2024

Received in revised form

October 30, 2024

Accepted December 17, 2024

Available online

December 17, 2024

Keywords:

Cycloaddition

Polarity

Solvent effect

Electron transfer

ABSTRACT

The correlation between coefficients from Dimroth equations and descriptors of global electron density transfer was explored based on the data available in the recent literature. We established that the obtained results should be very usable for the interpretation of the organic reactivity and molecular mechanisms.

© 2025 by the authors; licensee Growing Science, Canada.

1. Introduction


Cycloaddition processes are the most universal way for the preparation of many, different-type of carbo- and heterocyclic skeletons^{1–5}. Mechanistic aspects of the [4+2]- π -electron cycloaddition reactions (with special participations of (3+2) and (4+2) processes) still pay the great attention of many scientists^{6–10}. Even a few decades ago, the dogma about the concerted reorganization of the electron density, and synchronous formation of new sigma-bonds was treated as the most important fundament of the cycloaddition theory¹¹. This view was however rapidly undermined after historical publications by Huisgen and Mloston at the 1986^{12,13}. At this moment, the general classification of cycloaddition mechanisms include two areas: polar and non-polar processes⁹. Within the first group, several scenarios are possible, such as polar, synchronous mechanisms, polar asynchronous (one step – two stage) mechanisms and stepwise mechanisms with the zwitterionic intermediate^{14–17}. Alternatively, non-polar cycloadditions can be classified as polar, synchronous, non-polar asynchronous (biradicaloid) mechanisms and stepwise mechanisms with the biradical intermediate^{18–20}.

For the interpretation of mechanistic aspects of cycloaddition processes, different type theoretical and experimental, kinetic techniques are dedicated. In the framework we decided to do a short analysis of possible correlation between experimentally obtained reaction constants in the Dimroth equations^{21–23} and theoretically calculated values of the global electron density transfer²⁴. Dimroth constant E_T is a parameter characterizing polarity of the solvent. It is based on pyridinium N-phenolbetaine **X** as a test substance owing to a significant displacement of a solvatochromic band. Value of the E_T can be calculated using equation (1), which is the frequency of the absorption maximum in cm^{-1} in the examined solvent.

$$E_T = 2.859 \cdot 10^{-3} \cdot \tilde{\nu} \left[\frac{\text{kcal}}{\text{mol}} \right]$$

* Corresponding author


E-mail address karolina.kula@pk.edu.pl (K. Kula)




Global Electron Density Transfer (GEDT) during transition state is a good measure of the polar character of reaction²⁵. GEDT takes place from nucleophile to electrophile, and can be obtained by sum of natural atomic charges of nucleophile fragments calculated through a natural population analysis (NPA)^{26,27}.

2. Results and discussion


Although, evidently most recent publications include the analysis of the solvent effects based on DFT computational results²⁸⁻³¹, in the literature are available some examples of experimental, kinetic study in this area. These results are especially interesting for the processes with the participation of NO₂-activated components^{32,33}. So, the (3+2) cycloaddition with the participation of the nitronium – fluorenone derivative (**1**) as the TAC and E-3,3,3-trichloro-1-nitroprop-1-ene (**2**) is realized at the r.t and is realized with full selectivity yielded respective 4-nitroisoxazolidine with almost quantitatively yield^{34,35}. The kinetic study regarding this transformation includes the analysis of the influence of the polarity solvent on the rate constants, as well as the estimation of activation enthalpy and entropy.




The similar package of kinetic measurements were collected for the (4+2) cycloaddition reaction between cyclopentadiene (**4**) and E-2-phenylnitroethene (**5**)³⁶. In this case, the substituent effect was also experimentally evaluated for analogous reaction with the participation of other 2-arylnitroethenes.



The next example of the comprehensive kinetic study is a cycloaddition of Z-C,N-diphenylnitronium (**7**) with mentioned above 3,3,3-trichloro-1-nitroprop-1-ene³⁷.



The same nitronium was also tested as a component of the (3+2] cycloaddition to the gem-chloronitroethene (**9**)³⁸. In this case, next to solvent effects, the substituent effects and the activation parameters were also estimated.



Lastly, in the recent literature, the kinetic study on the hetero Diels-Alder reaction of E-2-(4-carbomethoxyphenyl)-1-cyano-1-nitroethene (**11**) and ethyl-vinyl ether (**12**) are available^{39,40}. This analysis was supported by estimation of the analysis of the substituent effects and the activation parameters.

Fortunately, for all mentioned processes, respective DFT mechanistic studies are additionally performed. So, it is possible to estimate the value of the electron density transfer between substructures. These descriptors, and experimental measured constants in Dimroth correlations are collected in **Table 1**.

Table 1. Key kinetic parameters for model cycloaddition processes

Reaction	Global electron density transfer (GEDT) [e]	Coefficient in the Dimroth equation
1+2	0.18	0.008
4+5	0.21	0.009
7+2	0.21	0.011
7+9	0.32	0.031
11+12	0.40	0.049

The correlation analysis exhibits the excellent^{41,42} relationship ($R > 0.99$) between coefficients from Dimroth equations and GEDT values.

$$\text{GEDT} = 5.13 \cdot E_T + 0.15 \quad (R=0.994)$$

So, in the evaluated range of Dimroth coefficients, this relation can be treated as general and usable for the prediction of the solvent effect based on theoretical estimated GEDTs.




Fig. 1. The graphical illustration of the correlation between coefficients from Dimroth equations and GEDT values.

3. Conclusion

Analysis of literature source exhibits, that independently of many theoretical works, in the recent time some valuable papers regarding the experimental kinetic studies were performed. Based on this information we decided to explore the theoretical background for the observed solvent effects. Our short analysis confirms without any doubts the existence of the correlation between coefficients from Dimroth equations and descriptors of global electron density transfer. In our opinion

this tool can be a simple and usable approach for the predictions of the organic reactivity and molecular mechanisms within cycloaddition reactions.

Grants and funding

This research was funded in whole by the National Science Centre, Poland, Miniatura 8 (grant number: 2024/08/X/ST4/00933).

References

- (1) Prakash, C., Singh, R. (2024) Synthesis of Fluorinated Six-Membered Nitrogen Heterocycles Using Microwave Irradiation. *Chem. Heterocycl. Compd.*, 60 (5–6), 216–229. <https://doi.org/10.1007/s10593-024-03323-1>.
- (2) Zhilitskaya, L. V., Yarosh, N. O. (2024) The Synthesis of Salts of Five-Membered Heterocyclic Compounds Based on N-Containing Cations/Anions (Microreview). *Chem. Heterocycl. Compd.*, 60 (5–6), 230–232. <https://doi.org/10.1007/s10593-024-03324-0>.
- (3) Varala, R., Kurra, M., Amanullah, M., Hussien, M., Alam, M. M. (2024,) Recent Methods in the Synthesis of Chromeno[2,3-d]Pyrimidines. *Chem. Heterocycl. Compd.* 60 (3–4), 111–117. <https://doi.org/10.1007/s10593-024-03302-6>.
- (4) Dotsenko, V. V., Sinotsko, A. E. (2024) The Synthesis and Properties of [1,2] Dithiopyridine Derivatives (Microreview). *Chem. Heterocycl. Compd.*, 60 (1–2), 32–34. <https://doi.org/10.1007/s10593-024-03288-1>.
- (5) Chaban, T. I., Lelyukh, M. I., Chaban, I. H., Kasyanchuk, O. Y. (2024) Approaches to the Synthesis of Thiazolo[3,2-a]Pyridines (Microreview). *Chem. Heterocycl. Compd.*, 60 (3–4), 124–126. <https://doi.org/10.1007/s10593-024-03305-3>.
- (6) Jasiński, R. (2021) On the Question of Stepwise [4+2] Cycloaddition Reactions and Their Stereochemical Aspects. *Symmetry (Basel)*, 13 (10), 1911. <https://doi.org/10.3390/sym13101911>.
- (7) Sadowski, M., Kula, K. (2024) Nitro-Functionalized Analogues of 1,3-Butadiene: An Overview of Characteristic, Synthesis, Chemical Transformations and Biological Activity. *Curr. Chem. Lett.*, 13 (1), 15–30. <https://doi.org/10.5267/j.ccl.2023.9.003>.
- (8) Jasiński, R., Dresler, E. (2020) On the Question of Zwitterionic Intermediates in the [3+2] Cycloaddition Reactions: A Critical Review. *Organics*, 1 (1), 49–69. <https://doi.org/10.3390/org1010005>.
- (9) Domingo, L. R., Sáez, J. A. (2009) Understanding the Mechanism of Polar Diels–Alder Reactions. *Org. Biomol. Chem.*, 7 (17), 3576. <https://doi.org/10.1039/b909611f>.
- (10) Basavanna, V., Puttappa, S., Chandramouli, M., Ningaiah, S. (2023,) Green Synthetic Methods for the Cycloaddition Reactions: A Mini Review. *Polycycl. Aromat. Compd.* 43 (10), 9377–9398. <https://doi.org/10.1080/10406638.2022.2162933>.
- (11) Huisgen, R. (1963) 1,3-Dipolar Cycloadditions. Past and Future. *Angew. Chemie Int. Ed. English*, 2 (10), 565–598. <https://doi.org/10.1002/anie.196305651>.
- (12) Huisgen, R., Mloston, G., Langhals, E. (1986) The First Two-Step 1,3-Dipolar Cycloadditions: Non-Stereospecificity. *J. Am. Chem. Soc.*, 108 (20), 6401–6402. <https://doi.org/10.1021/ja00280a053>.
- (13) Huisgen, R., Mloston, G., Langhals, E. (1986) The First Two-Step 1,3-Dipolar Cycloadditions: Interception of Intermediate. *J. Org. Chem.*, 51 (21), 4085–4087. <https://doi.org/10.1021/jo00371a039>.
- (14) Woliński, P., Kačka-Zych, A., Wróblewska, A., Wielgus, E., Dolot, R., Jasiński, R. (2023) Fully Selective Synthesis of Spirocyclic-1,2-Oxazine N-Oxides via Non-Catalysed Hetero Diels–Alder Reactions with the Participation of Cyanofunctionalised Conjugated Nitroalkenes. *Molecules*, 28 (12), 4586. <https://doi.org/10.3390/molecules28124586>.
- (15) Kula, K., Łapczuk, A., Sadowski, M., Kras, J., Zawadzińska, K., Demchuk, O. M., Gaurav, G. K., Wróblewska, A., Jasiński, R. (2022,) On the Question of the Formation of Nitro-Functionalized 2,4-Pyrazole Analogs on the Basis of Nitylimine Molecular Systems and 3,3,3-Trichloro-1-Nitroprop-1-Ene. *Molecules* 27 (23), 8409. <https://doi.org/10.3390/molecules27238409>.
- (16) Kula, K., Kačka-Zych, A., Łapczuk-Krygier, A., Jasiński, R. (2021) Analysis of the Possibility and Molecular Mechanism of Carbon Dioxide Consumption in the Diels–Alder Processes. *Pure Appl. Chem.*, 93 (4), 427–446. <https://doi.org/10.1515/pac-2020-1009>.
- (17) Fryźlewicz, A., Olszewska, A., Zawadzińska, K., Woliński, P., Kula, K., Kačka-Zych, A., Łapczuk-Krygier, A., Jasiński, R. (2022) On the Mechanism of the Synthesis of Nitrofunctionalised Δ^2 -Pyrazolines via [3+2] Cycloaddition Reactions between α -EWG-Activated Nitroethenes and Nitylimine TAC Systems. *Organics*, 3 (1), 59–76. <https://doi.org/10.3390/org3010004>.
- (18) Kačka-Zych, A., Pérez, P. (2021) Perfluorobicyclo[2.2.0]Hex-1(4)-Ene as Unique Partner for Diels–Alder Reactions with Benzene: A Density Functional Theory Study. *Theor. Chem. Acc.*, 140 (2), 17. <https://doi.org/10.1007/s00214-020-02709-6>.
- (19) Jasiński, R. (2020) A New Insight on the Molecular Mechanism of the Reaction between (Z)-C,N-Diphenylnitron and 1,2-Bismethylene-3,3,4,4,5,5-Hexamethylcyclopentane. *J. Mol. Graph. Model.*, 94. <https://doi.org/10.1016/j.jmgm.2019.107461>.
- (20) Jasiński, R. (2016) A Reexamination of the Molecular Mechanism of the Diels–Alder Reaction between

- Tetrafluoroethene and Cyclopentadiene. *React. Kinet. Mech. Catal.*, 119 (1), 49–57. <https://doi.org/10.1007/s11144-016-1038-1>.
- (21) Reichardt, C. (1965) Empirical Parameters of the Polarity of Solvents. *Angew. Chemie Int. Ed. English*, 4 (1), 29–40. <https://doi.org/10.1002/anie.196500291>.
- (22) Perrin, C. L., Agron, I., Bagno, A., Braslavsky, S. E., Fernandes, P. A., Gal, J.-F., Lloyd-Jones, G. C., Mayr, H., Murdoch, J. R., Nudelman, N. S., Radom, L., Rappoport, Z., Ruasse, M.-F., Siehl, H.-U., Takeuchi, Y., Tidwell, T. T., Uggerud, E., Williams, I. H. Dimroth–Reichardt Et Parameter. *IUPAC Standards Online*. De Gruyter. <https://doi.org/10.1515/iupac.94.0407>.
- (23) Tunuli, M. S., Rauf, M. A., Farhataziz. Dimroth's ET(30) as Parameters of Solvent Polarity: A Caveat. *J. Photochem.* 1984, 24 (4), 411–413. [https://doi.org/10.1016/0047-2670\(84\)80023-4](https://doi.org/10.1016/0047-2670(84)80023-4).
- (24) Domingo, L. R., Ríos-Gutiérrez, M. (2023) A Useful Classification of Organic Reactions Based on the Flux of the Electron Density. *Sci. Radices*, 2 (1), 1–24. <https://doi.org/10.58332/scirad2023v2i1a01>.
- (25) Domingo, L. R. (2016) Molecular Electron Density Theory: A Modern View of Reactivity in Organic Chemistry. *Molecules*, 21 (10), 1319. <https://doi.org/10.3390/molecules21101319>.
- (26) Reed, A. E. (1985) Weinstock, R. B., Weinhold, F. Natural Population Analysis. *J. Chem. Phys.*, 83 (2), 735–746. <https://doi.org/10.1063/1.449486>.
- (27) Domingo, L. R. (2014) A New C–C Bond Formation Model Based on the Quantum Chemical Topology of Electron Density. *RSC Adv.*, 4 (61), 32415–32428. <https://doi.org/10.1039/C4RA04280H>.
- (28) Dresler, E., Woliński, P., Wróblewska, A., Jasiński, R. (2023) On the Question of Zwitterionic Intermediates in the [3+2] Cycloaddition Reactions between Aryl Azides and Ethyl Propiolate. *Molecules*, 28 (24), 8152. <https://doi.org/10.3390/molecules28248152>.
- (29) Sadowski, M., Dresler, E., Zawadzińska, K., Wróblewska, A., Jasiński, R. (2024) Syn-Propanethial S-Oxide as an Available Natural Building Block for the Preparation of Nitro-Functionalized, Sulfur-Containing Five-Membered Heterocycles: An MEDT Study. *Molecules*, 29 (20), 4892. <https://doi.org/10.3390/molecules29204892>.
- (30) Zawadzińska, K., Kula, K. (2021) Application of β -Phosphorylated Nitroethenes in [3+2] Cycloaddition Reactions Involving Benzonitrile N-Oxide in the Light of a DFT Computational Study. *Organics*, 2 (1), 26–37. <https://doi.org/10.3390/org2010003>.
- (31) Łapczuk-Krygier, A., Jaśkowska, J., Jasiński, R. (2018) The Influence of Lewis Acid Catalyst on the Kinetic and Molecular Mechanism of Nitrous Acid Elimination from 5-Nitro-3-Phenyl-4,5-Dihydroisoxazole: DFT Computational Study. *Chem. Heterocycl. Compd.*, 54 (12), 1172–1174. <https://doi.org/10.1007/s10593-019-02410-y>.
- (32) Sadowski, M., Synkiewicz-Musialska, B., Kula, K. (2024.) (1E,3E)-1,4-Dinitro-1,3-Butadiene—Synthesis, Spectral Characteristics and Computational Study Based on MEDT, ADME and PASS Simulation. *Molecules* 29 (2), 542. <https://doi.org/10.3390/molecules29020542>.
- (33) Sadowski, M., Kula, K. (2024) Unexpected Course of Reaction Between (1E,3E)-1,4-Dinitro-1,3-Butadiene and N-Methyl Azomethine Ylide—A Comprehensive Experimental and Quantum-Chemical Study. *Molecules*, 29 (21), 5066. <https://doi.org/10.3390/molecules29215066>.
- (34) Jasiński, R., Mróz, K. (2015.) Kinetic Aspects of [3+2] Cycloaddition Reactions between (E)-3,3,3-Trichloro-1-Nitroprop-1-ene and Ketonitrones. *React. Kinet. Mech. Catal.* 116 (1), 35–41. <https://doi.org/10.1007/s11144-015-0882-8>.
- (35) Jasiński, R., Mróz, K., Kačka, (2016) A. Experimental and Theoretical DFT Study on Synthesis of Sterically Crowded 2,3,3,(4)5-Tetrasubstituted-4-Nitroisoxazolidines via 1,3-Dipolar Cycloaddition Reactions Between Ketonitrones and Conjugated Nitroalkenes. *J. Heterocycl. Chem.*, 53 (5), 1424–1429. <https://doi.org/10.1002/jhet.2442>.
- (36) Kwiatkowska, M. (2008) Stereochemiczne i Kinetyczne Studia [2+4] Cykloaddycji Cyklopentadienu Ze Sprzężonymi Nitroalkenami, Cracow University of Technology, PhD Thesis,.
- (37) Jasiński, R. (2004) Reakcje [2+3] Cykloaddycji Z-C,N-Difenylonitronu i C,C,N-Trifenylonitronu Ze Sprzężonymi Nitroalkenami, Politechnika Krakowska. PhD Thesis,
- (38) Mikulska, M. (2015) Teoretyczne i Eksperymentalne Studia Reakcji [2+3] Cykloaddycji 1–Podstawionych Nitroetenów z (Z)–C,N–Diarylonitronami Oraz Termoliza Uzyskanych Adduktów, Cracow University of Technology, PhD ThesisKula.
- (39) Jasiński, R., Kubik, M., Łapczuk-Krygier, A., Kačka, A., Dresler, E., (2014) Boguszewska-Czubara, A. An Experimental and Theoretical Study of the Hetero Diels–Alder Reactions between (E)-2-Aryl-1-Cyano-1-Nitroethenes and Ethyl Vinyl Ether: One-Step or Zwitterionic, Two-Step Mechanism? *React. Kinet. Mech. Catal.*, 113 (2), 333–345. <https://doi.org/10.1007/s11144-014-0753-8>.
- (40) Fringuelli, F., Matteucci, M., Piermatti, O., Pizzo, F., Burla, M. C. (2001) [4 + 2] Cycloadditions of Nitroalkenes in Water. Highly Asymmetric Synthesis of Functionalized Nitronates. *J. Org. Chem.*, 66 (13), 4661–4666. <https://doi.org/10.1021/jo010182v>.
- (41) Schwetlick, K. *Kinetische Methoden Zur Untersuchung von Reaktionsmechanismen*, VEB Deutscher Verlag der Wissenschaften: Berlin, 1971.
- (42) Reichardt, C., Welton, T. (2010) *Solvents and Solvent Effects in Organic Chemistry*, Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim. <https://doi.org/10.1002/9783527632220>.



© 2025 by the authors; licensee Growing Science, Canada. This is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC-BY) license (<http://creativecommons.org/licenses/by/4.0/>).